

How to give the right amount of ACETAMINOPHEN (also known as Tylenol) is different depending on which medicine you plan to give.

Dose: Give every 4 to 6 hours as needed for fever or pain. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**
Do **NOT** use with any other medicine containing acetaminophen.



Weight →	6 to 11 pounds (3 to 5 kilograms)	12 to 17 pounds (5 to 7 kilograms)	18 to 23 pounds (8 to 10 kilograms)	24 to 35 pounds (11 to 15 kilograms)
Age →	0 to 3 months	4 to 11 months	12 to 23 months	2 to 3 years
Infant's Acetaminophen (160 mg / 5 mL)	1.25 mL 	2.5 mL 	3.75 mL 	_____
Children's Acetaminophen (160 mg / 5 mL)	_____	_____	_____	5 mL 
Children's Acetaminophen Chewables (160 mg)	_____	_____	_____	1 tablet 
Children's Acetaminophen Dissolvable Packets (160 mg / powder pack)	_____	_____	_____	_____
Adult's Acetaminophen Tablets (325 mg)	_____	_____	_____	_____
Adult's Acetaminophen Tablets (500 mg)	_____	_____	_____	_____

How to give the right amount of IBUPROFEN (also known as Motrin, Advil) is different depending on which type of ibuprofen you plan to give.

Dose: Give every 6 hours if needed, for fever or pain. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**
Do **NOT** use with any other medicine containing ibuprofen.

















Weight	Age	Infant's Ibuprofen Drops (50 mg / 1.25 mL)	Children's Liquid Ibuprofen (100 mg / 5 mL)	Children's Ibuprofen Chewable Tablets (50 mg)	Junior Strength Ibuprofen Tablets (100 mg)	Adult's Ibuprofen Tablets (200 mg)
0 to 11 pounds (up to 5 kilograms)	0 to 5 months	_____	_____	_____	_____	_____
12 to 17 pounds (about 6 to 7 kilograms)	6 to 11 months	1.25 mL 	2.5 mL 	_____	_____	_____
18 to 23 pounds (about 8 to 10 kilograms)	12 to 23 months	1.875 mL 	4 mL 	_____	_____	_____
24 to 35 pounds (about 11 to 15 kilograms)	2 to 3 years	2.5 mL 	5 mL → 	2 tablets 	_____	_____
36 to 47 pounds (about 16 to 21 kilograms)	4 to 5 years	3.75 mL 	 ← 7.5 mL	3 tablets 	_____	_____
48 to 59 pounds (about 22 to 26 kilograms)	6 to 8 years	5 mL 	10 mL → 	4 tablets 	2 tablets 	1 tablets 
60 to 71 pounds (about 27 to 32 kilograms)	9 to 10 years	_____	 ← 12.5 mL	5 tablets 	2 ½ tablets 	1 tablet 
72 to 95 pounds (about 33 to 43 kilograms)	11 years	_____	 ← 15 mL	6 tablets 	3 tablets 	1 ½ tablets 
96 pounds or more (44 kilograms or more)	12 years or older	_____	10 mL + 10 mL = 20 mL 10 mL →  and 10 mL → 	8 tablets 	4 tablets 	2 tablets 

How to give the right amount of DIPHENHYDRAMINE (also known as Benadryl) is different depending on which type of Diphenhydramine you plan to give.

Dose: Give every 6 hours if needed. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**
Do **NOT** use with any other medicine with diphenhydramine in it.



Weight →	20 to 24 pounds (about 9 to 10 kilograms)	25 to 37 pounds (about 11 to 16 kilograms)	38 to 49 pounds (about 17 to 22 kilograms)	50 to 99 pounds (about 23 to 45 kilograms)	100 pounds or more (46 kilograms or more)
Children's Liquid Diphenhydramine (12.5 mg / 5 mL)	4 mL 	5 mL 	7.5 mL 	10 mL 	—
Children's Diphenhydramine Chewable Tablets (12.5 mg)	—	1 tablet 	1 ½ tablets 	2 tablets 	4 tablets 
Diphenhydramine Tablets (25 mg)	—	½ tablet 	½ tablet 	1 tablet 	2 tablets 
Diphenhydramine Capsules (25 mg)	—	—	—	1 capsule 	2 capsules 

Do not give to children less than 2 years of age.

Do not give to children 2 to 6 years of age unless your doctor tells you to.